

Expansivity properties and rigidity for non-recurrent exponential maps

Anna Miriam Benini *

CRM, Scuola Normale Superiore

Piazza Cavalieri 7

Pisa, Italy

ambenini@gmail.com

October 5, 2012

Abstract

We show that an exponential map $f_c(z) = e^z + c$ whose singular value c is combinatorially non-recurrent and non-escaping is uniquely determined by its combinatorics, i.e. the pattern in which its dynamic rays land together. We do this by constructing puzzles and parapuzzles in the exponential family. We also prove a theorem about hyperbolicity of the postsingular set in the case that the singular value is non-recurrent. Finally, we show that boundedness of the postsingular set implies combinatorial non-recurrence if c is in the Julia set.

Introduction

Consider the family of exponential maps $f_c(z) = e^z + c$, with $c \in \mathbb{C}$. It is a natural and relevant question to ask whether there exists a dynamical condition under which two maps in this family are conformally conjugate, and hence correspond to the same parameter up to translation by $2\pi i$. In this paper we give a sufficient combinatorial condition for this to happen, under the hypothesis of a specific kind of non-recurrence.

The set of *escaping points*

$$I(f_c) := \{z \in \mathbb{C} : |f_c^n(z)| \rightarrow \infty\}$$

*This work was partly supported by a DGAPA-UNAM fellowship at IMATE, Cuernavaca, Mexico, as well as by the Institute for Mathematical Sciences in Stony Brook, US.

can be naturally described as a union of injective curves, called *dynamic rays* or *hairs*, labelled by sequences in $\mathbb{Z}^{\mathbb{N}}$ in such a way that the dynamics of f_c on the rays is conjugate to the dynamics of the shift map σ on $\mathbb{Z}^{\mathbb{N}}$. We refer the reader to [BDHRGH1], [SZ1] and Section 2.1 in this paper for existence and properties of dynamic rays in the exponential family. Informally, dynamic rays can be thought of as curves from $(0, \infty)$ to \mathbb{C} , which tend to infinity as the parameter $t \rightarrow \infty$. On the other side, the behavior as $t \rightarrow 0$ can be far more complicated; if the limit exists, we say that the dynamic ray *lands*. A dynamic ray is called *periodic* or *preperiodic* if it is a periodic or preperiodic set under the dynamic of f_c ; it is shown in [Re1] that periodic and preperiodic rays land unless their forward orbit contains the singular value.

Two periodic or preperiodic dynamic rays landing at the same point, together with their common endpoint, form a curve Γ disconnecting the plane. Given two such dynamic rays, we say that two points are *separated* (by Γ) if they belong to different connected components of $\mathbb{C} \setminus \Gamma$.

A fundamental question is under which conditions the set of periodic and preperiodic rays landing together, which is just a combinatorial data, completely encodes the actual dynamics, and hence determines uniquely the position of the map in parameter plane. This property is usually referred to as *rigidity*. Stated in this terms, rigidity can be asked for parameters which do not belong to the closure of hyperbolic components (or equivalently, for which all periodic point are repelling). Two maps f_c, f_c are called *combinatorially equivalent* if their periodic and preperiodic rays land together in the same pattern (see Section 4.2 for a more precise definition).

A positive answer to the question of rigidity would imply the famous *density of hyperbolicity conjecture*, one of the main open problems in one-dimensional complex dynamics. Conjecturally, hyperbolic maps are dense in appropriate spaces of functions, like in the space of unicritical polynomials $P_c^D(z) = z^D + c$ and in the space of exponential maps $f_c(z) = e^z + c$. The question of rigidity is also strictly related to the MLC conjecture, according to which the Mandelbrot set is locally connected. See [RS3] for a paper making a parallel between rigidity problems in exponential versus polynomial setting.

In this paper we study exponential maps whose singular value is *non-recurrent*, that is the parameters c for which the postsingular set

$$\mathcal{P}(f_c) = \overline{\bigcup_{n>0} \{f_c^n(c)\}}$$

does not contain the singular value c itself. Observe that for an exponential map f_c , boundedness of $\mathcal{P}(f_c)$ implies non-recurrence.

The first result in this paper is hyperbolicity of $\mathcal{P}(f_c)$ in the case that c is non-recurrent. This is a weak version of a theorem of Mañé for non-recurrent exponential maps with unbounded postsingular set (see [Ma], [ST] for rational maps and [RvS] for

a wider generalization in the transcendental setting; see also [GKS] for meromorphic functions with bounded postsingular set). As a consequence of the aforementioned Mañé's Theorem, the boundary of Siegel disks is contained in the ω -limit set of a recurrent singular value, hence in the exponential family non-recurrence of c implies absence of Siegel Disks (see [RvS, Corollary 2.10]).

A forward invariant set K is *hyperbolic* (with respect to the Euclidean metric) if there exists $\bar{k}, \eta > 1$ such that for any $k > \bar{k}$ and for any $z \in K$, $|(f^k)'(z)| > \eta$.

Theorem A (Hyperbolicity of the postsingular set). *Let $f_c(z) = e^z + c$ such that $c \in J(f_c)$ is non-recurrent. Then $\mathcal{P}(f_c)$ is hyperbolic.*

The rest of the paper is devoted to problems related to rigidity. We first define Yoccoz puzzle and parapuzzles in the exponential setting. We then restrict ourselves to the class of *combinatorially non-recurrent* parameters: a parameter c is *combinatorially non-recurrent* if there is a suitable collection (see Section 4) of preperiodic rays which separate the singular value from the postsingular set. Under the assumption of combinatorial non-recurrence we prove that two combinatorially equivalent maps are quasiconformally conjugate.

Theorem B. *Let c, c' be non-escaping parameters, and f_c be combinatorially non-recurrent. If $f_{c'}$ is combinatorially equivalent to f_c , then $f_{c'}$ is quasiconformally conjugate to f_c .*

In Section 6 we show actual rigidity for combinatorially non-recurrent parameters. For parameters with bounded postsingular set, this can be obtained as a corollary of Theorem B together with a result of [RvS] on the absence of invariant line fields (see also [MS]). Otherwise, the proof is more involved. The final result is the following:

Theorem C. *Let c, c' be non-escaping parameters, and f_c be combinatorially non-recurrent. If $f_{c'}$ is combinatorially equivalent to f_c , then $c' = c$.*

We conclude by showing that if $c \in J(f)$, boundedness of the post-singular set is a sufficient condition to imply combinatorial non-recurrence (see Section 7).

Theorem D. *Let f_c be an exponential map such that $c \in J(f_c)$ and $\mathcal{P}(f_c)$ is bounded. Then c is combinatorially non-recurrent.*

The condition $c \in J(f_c)$ is in order to exclude hyperbolic and parabolic parameters, which are non-recurrent in the classical sense but not in the combinatorial sense. The proof of Theorem D is independent from the rest of the paper, and relies on one side on known rigidity results for polynomials, and on the other side on the combinatorial similarity between exponentials and unicritical polynomials. It also uses a theorem from [BL] about accessibility of the singular value.

Acknowledgements

I am gratefully indebted to my advisor Misha Lyubich for suggesting this problem and for fundamental discussions about this topic. I am also thankful to Peter Makienko, Carsten Petersen, Lasse Rempe, Dierk Schleicher and Anna Zdunik for related discussions.

Notation and terminology

The complex plane is \mathbb{C} , the open unit disk is \mathbb{D} and the Riemann sphere is $\hat{\mathbb{C}}$. We denote by $D_r(z)$ a disk of radius r centered at the point z . By a neighborhood of a point we mean a connected, simply connected neighborhood.

The Euclidean diameter of a set U' is denoted by $\text{diam } U'$, while the Euclidean length of a curve γ is denoted by $\ell(\gamma)$. If U admits a normalized hyperbolic metric, the hyperbolic diameter of a set $U' \subset U$ is denoted by $\text{diam}_U(U')$, while the hyperbolic length of a curve $\gamma \subset U$ is denoted by $\ell_U(\gamma)$.

The Julia set of an exponential map f is denoted by $J(f)$, and its Fatou set by $F(f)$. A parameter c is called *non-escaping* if $f_c^n(c) \not\rightarrow \infty$. It is called *hyperbolic* if it has an attracting periodic orbit, *parabolic* if it has an indifferent periodic orbit with rational multiplier, and *Misiurewicz* if the orbit of c is finite. It is called *Siegel* (respectively *Cremer*) if it has an indifferent periodic orbit $\{z_i\}_{i=1}^q$ with irrational multiplier in a neighborhood of which f_c^q is linearizable (respectively non-linearizable). It is called *non-recurrent* if the singular value is non-recurrent. A maximal open set of parameters which are hyperbolic is called a *hyperbolic component*. Parabolic, Siegel and Cremer parameters are on the boundaries of hyperbolic components.

For an exponential map f and a topological disks V , we call $\mathcal{C}(n, V)$ the set of all connected components of $f^{-n}(V)$, and $\mathcal{C}_P(n, V)$ the set of connected components of $f^{-n}(V)$ which intersect $\mathcal{P}(f)$.

1 Quasiconformal maps and holomorphic motions

In this section we recall the definition and the main properties of quasiconformal maps and of holomorphic motions. These results and their proof can be found for example in [Ah]; they will be used in Sections 4,5,6.

Let $\psi : \mathbb{C} \rightarrow \mathbb{C}$ be a homeomorphism with partial derivatives $\partial_z \psi, \partial_{\bar{z}} \psi$ in the sense of distribution; we associate to ψ its *Beltrami coefficient* $\mu_\psi := \frac{\partial_{\bar{z}} \psi}{\partial_z \psi}$. By definition $\mu_\psi(z)$ is a measurable function. The map ψ is called *quasiconformal* if $\|\mu_\psi\|_\infty < 1$. The quantity $K := \sup_z \frac{\mu_\psi(z)+1}{1-\mu_\psi(z)}$ is called the *dilatation* of ψ ; if ψ has dilatation K , it is called K -quasiconformal. Inverses of K -quasiconformal maps are K -quasiconformal; moreover, if g is conformal and ψ is K -quasiconformal, $g \circ \psi$ as well as $\psi \circ g$ are

K -quasiconformal.

Lemma 1.1 (Weyl's Lemma). *If f is a quasiconformal map with $\mu_f = 0$ outside a set of Lebesgue measure zero, then f is conformal.*

We say that a sequence of functions $\{\psi_n\}$ has a limit function ψ if there is a subsequence converging to ψ uniformly on compact sets. One of the remarkable properties of K -quasiconformal maps is their compactness.

Lemma 1.2. *Let $\{\psi_n\}$ be a family of K -quasiconformal functions which coincide on at least three points. Then there exists at least one limit function ψ , and it is K -quasiconformal.*

By definition, a quasiconformal map ψ induces a Beltrami coefficient with $\|\mu_\psi\|_\infty < 1$; it is a natural question to ask whether any allowable Beltrami coefficient (which can be thought of as a measurable function with modulus less than 1) corresponds to a quasiconformal map. The answer is provided by the Measurable Riemann Mapping Theorem.

Theorem 1.3 (Measurable Riemann Mapping Theorem). *Let $\{\mu_\lambda\}$, $\|\mu_\lambda\|_\infty < 1$, be a family of Beltrami coefficients depending holomorphically on a parameter λ . Then there is a family of quasiconformal maps ψ_λ , depending holomorphically on λ , such that $\mu_\lambda = \partial_z \psi_\lambda / \partial_{\bar{z}} \psi_\lambda$.*

A Beltrami coefficient defines a measurable field of ellipses in the tangent space, with bounded ratio between the length of the major and minor axis. On the other side, a measurable field of ellipses with bounded ratio between the major and minor axis defines a Beltrami coefficient. This field of ellipses defines a so-called *conformal structure* on \mathbb{C} . For a holomorphic map g , $\mu_g = 0$, and the induced field of ellipses is in fact a field of circles. We refer to this field of circles as the *standard conformal structure* σ_0 on \mathbb{C} .

Given a holomorphic function f , an *invariant line field* is a field of lines defined in the tangent space which is invariant under f . A Beltrami coefficient μ defines an invariant line field through the directions of the major axis of the induced field of ellipses in the tangent space. The induced line field is invariant if and only if $f^* \mu = \mu$. The following theorem is a special case of Theorem 1.1 in [RvS].

Theorem 1.4 (Absence of invariant line fields). *Let $f_c(z) = e^z + c$ such that $\mathcal{P}(f_c)$ is bounded. Then $J(f_c)$ supports no invariant line fields.*

Holomorphic motions are another widely used tool in one-dimensional complex dynamics.

Definition 1.5. Let $(\Lambda, *)$ be a topological disk in \mathbb{C} with a marked point $*$, and let X be a subset of the Riemann sphere. A *holomorphic motion* \mathbf{h} of X over $(\Lambda, *)$ is a

family of injections $h_\lambda : X \rightarrow \mathbb{C}$, $\lambda \in \Lambda$, depending holomorphically on λ for each fixed $x \in X$, and such that h_* is the identity. Define $X_\lambda := h_\lambda(X)$. A set X is said to *move holomorphically* over Λ if such a holomorphic motion exists.

The following theorem, due to [MSS] and improved by [SI], collects the most relevant results about holomorphic motions. See Theorem 1.3 and Lemma 1.1 in [SI].

Theorem 1.6 (Lambda Lemma). *Let \mathbf{h} be a holomorphic motion of a set $X \subset \mathbb{C}$ over the disk $(\Lambda, *)$. Then the two following facts hold:*

- \mathbf{h} extends to a continuous holomorphic motion of \mathbb{C} over $(\Lambda, *)$;
- The maps h_λ are K -quasiconformal with dilatation K depending only on the hyperbolic distance between λ and $*$. The constant K tends to infinity as $\lambda \rightarrow \partial\Lambda$.

2 Background in exponential dynamics

In this section we recollect the relevant information needed about the exponential family. In particular, we introduce dynamic and parameter rays and we describe the structure that they induce in the dynamical and parameter plane respectively. Both have been first introduced in [BDHRGH1] with the term *hairs*, and have been fully classified in [SZ1] and [FS].

2.1 Dynamic rays

For all of this section, $f_c(z) = e^z + c$. Let $F(t) = e^t - 1$ be a model function for real exponential growth, σ be the left-sided shift map acting on $\mathbb{Z}^{\mathbb{N}}$.

An *exponentially bounded address*, or just *address* is a sequence $\mathbf{s} = s_0 s_1 \dots \in \mathbb{Z}^{\mathbb{N}}$, such that $|s_i| \leq F^i(x)$ for some $x \in \mathbb{R}$ and for every $i \in \mathbb{N}$. The set of addresses is denoted by \mathcal{S} . An address is called *periodic* or *preperiodic* if it is a periodic or preperiodic sequence.

For any two sequences $\mathbf{s} = s_0 s_1 s_2 \dots$ and $\mathbf{s}' = s'_0 s'_1 s'_2 \dots$ in \mathcal{S} , we consider the distance

$$|\mathbf{s} - \mathbf{s}'| = \sum_{s_i \neq s'_i} \frac{1}{2^i}. \quad (2.1)$$

An address is called *bounded* if $\|\mathbf{s}\| := \sup_i |s_i| < \infty$. We refer to \mathcal{S} as the *combinatorial space* for f . Observe that \mathcal{S} is endowed with the lexicographic order.

Definition, existence and properties of *dynamic rays* for the exponential family are summarized in the following theorem ([SZ1], Proposition 3.2 and Theorem 4.2):

Theorem 2.1 (Dynamic rays, [SZ1]). *Let $c \in \mathbb{C}$ and $\mathbf{s} \in \mathcal{S}$ be an exponentially bounded address. Then, there exist $t_{\mathbf{s},c}$ and a unique maximal injective curve $g_{\mathbf{s}}^c : (t_{\mathbf{s},c}, \infty) \rightarrow I(f_c)$ such that*

$$(a) \quad f_c(g_{\mathbf{s}}^c(t)) = g_{\sigma_{\mathbf{s}}}^c(F(t)).$$

$$(b) \quad f_c^n(g_{\mathbf{s}}^c(t)) = 2\pi i s_n + F^n(t) + o(e^{-F^n(t)}) \text{ as } t \rightarrow \infty.$$

Moreover,

$$(c) \quad \text{For any } \mathbf{s}, g_{\mathbf{s}}^c(t) \text{ depends analytically on } c \text{ unless } c \in g_{\sigma^n \mathbf{s}}^c \text{ for some } n \in \mathbb{N}.$$

The curve $g_{\mathbf{s}}^c$ is called the *dynamic ray* (or just *ray*) of address \mathbf{s} . If c is non-escaping, $t_{\mathbf{s},c}$ does not depend on c . We omit the index c and simply write $g_{\mathbf{s}}(t)$ when this creates no ambiguity. The estimates in (b) together with the fact that dynamic rays do not intersect induce a *vertical order* on dynamic rays near infinity, corresponding to the lexicographic order on \mathcal{S} . Injectivity together with (c) imply that the functions $g_{\mathbf{s}}^c \circ (g_{\tilde{\mathbf{s}}}^c)^{-1}$ define a holomorphic motion of the dynamic ray $g_{\mathbf{s}}^c$ over any neighborhood of \tilde{c} on which the ray $g_{\tilde{\mathbf{s}}}^c(t)$ is well defined.

A dynamic ray $g_{\mathbf{s}}^c$ is called *periodic* (resp. *preperiodic*) if \mathbf{s} is a periodic sequence (resp. strictly preperiodic). A dynamic ray $g_{\mathbf{s}}^c$ is said to *land* at $z \in \mathbb{C}$ if $\lim_{t \rightarrow t_{\mathbf{s},c}} g_{\mathbf{s}}^c(t) = z$. It is shown in [Re1] that periodic and preperiodic dynamic rays land unless one of their forward images contains the singular value.

Given a set \mathcal{A} of addresses, we say that the rays with addresses in \mathcal{A} for $f_c, f_{c'}$ *land together in the same pattern* whenever the following condition is satisfied: two rays $g_{\mathbf{s}}^c, g_{\mathbf{s}'}^{c'}$ with addresses in \mathcal{A} land together if and only if $g_{\mathbf{s}}^{c'}, g_{\mathbf{s}'}^{c'}$ land together. If all periodic and preperiodic dynamic rays for f_c and $f_{c'}$ land together in the same pattern, we say that f_c and $f_{c'}$ are *combinatorially equivalent*.

We have the following result about dependence on the external address \mathbf{s} (see e.g. [Re2, Lemma 3.2]).

Lemma 2.2 (Transversal Continuity). *Let $\{\mathbf{s}_n\} \subset \mathcal{S}$ be a sequence of addresses converging to an address \mathbf{s} . Then $g_{\mathbf{s}_n}(t)$ converges uniformly to $g_{\mathbf{s}}(t)$ on all intervals $[t_*, \infty]$ with $t_* > t_{\mathbf{s},c}$.*

The following result is well known and its proof is provided for completeness.

Lemma 2.3 (Landing of preimages of rays). *Let $f(z) = e^z + c$ be an exponential map, and $\mathbf{s}_1 < \mathbf{s}_2$ be two addresses such that $g_{\mathbf{s}_1}$ and $g_{\mathbf{s}_2}$ land together at some point $z \neq c$. If $\overline{g_{\mathbf{s}_1} \cup g_{\mathbf{s}_2}}$ separates c from $-\infty$, for any $k \in \mathbb{Z}$, $g_{k\mathbf{s}_1}$ lands together with $g_{(k-1)\mathbf{s}_2}$. If not, for any $k \in \mathbb{Z}$, $g_{k\mathbf{s}_1}$ lands together with $g_{k\mathbf{s}_2}$.*

Proof. As f is a local homeomorphism near z , preimages of $g_{\mathbf{s}_1}$ and $g_{\mathbf{s}_2}$ land together pairwise at preimages of z . If the curve $\gamma := \overline{g_{\mathbf{s}_1} \cup g_{\mathbf{s}_2}}$ has winding number one with

respect to c , the imaginary part increases by 2π along any connected component of $f^{-1}(\gamma)$. If the winding number is zero, there is no increase in the imaginary part. By the asymptotic estimates in Theorem 2.1, any two rays in a connected component of $f^{-1}(\gamma)$ have to differ by exactly one unit in the first entry of their addresses in the first case, and by none in the second case. \square

2.2 Parameter rays, parabolic wakes, Misiurewicz wakes

The set of escaping parameters also consists of curves tending to infinity, called *parameter rays* (see [FS], Theorem 3.7).

Theorem 2.4 (Parameter rays). *Let $\mathbf{s} \in \mathcal{S}$. Then there is $t_{\mathbf{s}} > 0$, and a unique maximal injective curve $G_{\mathbf{s}} : (t_{\mathbf{s}}, \infty) \rightarrow \mathbb{C}$, such that, for all $t > t_{\mathbf{s}}$, $c = G_{\mathbf{s}}(t)$ if and only if $c = g_{\mathbf{s}}^c(t)$. Also, $|G_{\mathbf{s}}(t) - (t + 2\pi i s_0)| \rightarrow 0$ as $t \rightarrow \infty$.*

Recall that a hyperbolic component W is a maximal connected set of parameters with an attracting periodic orbit of the same period. There is a unique hyperbolic component W_0 of period 1 (the period of a hyperbolic component W is the period of the attracting periodic orbit for parameters in W). Parameter rays are approximated from above and from below by other parameter rays and by curves in hyperbolic components; see [BR, Lemma 7.1], as well as the proof of the Squeezing Lemma in [RS2, Section 4] and the proof that the exponential bifurcation locus is not locally connected in [RS3, Theorem 5].

The landing pattern of parameter rays with periodic and preperiodic addresses carves the structure of parameter plane, and is related to the landing pattern of dynamic rays in dynamical plane, as well as the position of hyperbolic components. This is exemplified in the next two theorems. The first theorem describes the relation between the landing pattern of parameter and dynamic rays with periodic address (see Proposition 4 and Proposition 5 in [Re1], as well as [RS1]).

Proposition 2.5 (Parabolic wakes). *Given a hyperbolic component W , there are exactly two parameter rays $G_{\mathbf{s}^+}, G_{\mathbf{s}^-}$ of periodic addresses \mathbf{s}^+ and \mathbf{s}^- which land together on ∂W and separate W from $-\infty$. The connected component $\mathcal{W}(W)$ of $\mathbb{C} \setminus \overline{G_{\mathbf{s}^+} \cup G_{\mathbf{s}^-}}$ containing W is called the (parabolic) wake of W . The dynamic rays $g_{\mathbf{s}^+}, g_{\mathbf{s}^-}$ and any of their forward iterates move holomorphically over $\mathcal{W}(W)$, and they land together in the dynamical plane for f_c if and only if c belongs to $\mathcal{W}(W)$. The rays $g_{\mathbf{s}^+}$ and $g_{\mathbf{s}^-}$ are called characteristic dynamic rays, while $G_{\mathbf{s}^+}, G_{\mathbf{s}^-}$ are called characteristic parameter rays.*

If W_1, W_2 are two hyperbolic components such that $\partial W_1 \cap \partial W_2 \neq \emptyset$, and the period of W_2 is higher than the period of W_1 , we say that we say that W_2 is *attached* to W_1 .

The situation for preperiodic addresses is slightly more complicated. However, it is shown in [LSV] (Theorem 3.4 and Corollary 3.5) that parameter rays with preperiodic

addresses land at Misiurewicz parameters, and that each Misiurewicz parameter c_0 is the landing point of finitely many parameter rays $G_{\mathbf{s}_1} \dots G_{\mathbf{s}_q}$ with preperiodic addresses. Moreover, the dynamical rays of addresses $\mathbf{s}_1 \dots \mathbf{s}_q$ land at c_0 in the dynamical plane for f_{c_0} . It is also not hard to prove that any Misiurewicz parameter is contained in a parabolic wake attached to W_0 , by using the fact that all repelling fixed points are landing points of periodic dynamic rays. Using these results and the fact that dynamic rays move holomorphically wherever they are well defined, we obtain the following.

Proposition 2.6 (Misiurewicz wakes). *Let c be a Misiurewicz parameter belonging to a parabolic wake \mathcal{W} attached to W_0 . Let $G_{\mathbf{s}_1} \dots G_{\mathbf{s}_q}$ be the parameter rays landing at c , $G_{\mathbf{s}^+}$ and $G_{\mathbf{s}^-}$ be the characteristic parameter rays bounding \mathcal{W} . The connected components of $\mathcal{W} \setminus \bigcup_i G_{\mathbf{s}_i}$ are called Misiurewicz wakes. The dynamic rays $g_{\mathbf{s}_1}^c \dots g_{\mathbf{s}_q}^c$ move holomorphically in each Misiurewicz wake, and they land together in the dynamical plane for c if and only if c belongs to one of the Misiurewicz wakes not containing $G_{\mathbf{s}^+}$ and $G_{\mathbf{s}^-}$ in its boundary.*

2.3 Fibers

The *extended (parameter) fiber* of a parameter c_0 with all periodic orbits repelling is the set of parameters which cannot be separated from c_0 by a pair of parameter rays with periodic or preperiodic addresses landing together. The *reduced (parameter) fiber* of a parameter c_0 is the extended fiber intersected with the set of non-escaping parameters. For parameters with indifferent or attracting periodic orbits, also curves formed only by parameter rays, hyperbolic parameters and finitely many parabolic parameters have to be considered as separation lines (see [RS3] for alternative definitions).

The *dynamical fiber* of a point z_0 is the set of points which cannot be separated from z_0 by a pair of dynamic rays with periodic or preperiodic addresses landing together.

A fiber is *trivial* if it contains at most one non-escaping parameter or point.

Fibers cannot contain the accumulation sets of arbitrary many parameter rays (see [[RS3], Lemma 17]):

Lemma 2.7. *Any extended fiber contains the accumulation set of at most finitely many parameter rays, and these parameter rays have bounded addresses.*

3 Hyperbolicity of the postsingular set

This section is dedicated to prove Theorem A, which will be used in the proof of Theorem B. The proof of Theorem A itself will not be used later. For all of this section, let $f(z) = e^z + c$ with $c \in J(f)$ non-recurrent. This two conditions imply that $J(f) = \mathbb{C}$. Recall that for a topological disk V , $\mathcal{C}(n, V)$ is the set of connected components of $f^{-n}(V)$, and $\mathcal{C}_P(n, V)$ is the set of connected components of $f^{-n}(V)$ which intersect

$\mathcal{P}(f)$. By forward invariance of $\mathcal{P}(f)$, if $U \in \mathcal{C}_P(n, V)$, $f^j(U) \in \mathcal{C}_P(n - j, V)$ for any $j \leq n$. Similarly, if $V' \subset V$, and $U' \in \mathcal{C}_P(n, V')$, then $U' \subset U$ for some $U \in \mathcal{C}_P(n, V)$.

We recall that a family of univalent functions $\{\varphi_k\}$, $\varphi_k : V \rightarrow \mathbb{C}$, with V simply connected, is *normal* if every sequence either has a convergent subsequence, or escapes any compact set (see [Mi]). By Montel's Theorem, any family omitting three values is normal. So, for any V simply connected neighborhood of a point $z \in \mathbb{C}$ which omits a periodic orbit of period > 3 , the family of univalent inverse branches of f defined on V is normal.

The following proposition is a special case of Theorem 2.7 in [RvS]. The proof in our case is inspired by [Ma] and much simpler, so we include it for completeness.

Proposition 3.1 (Univalent pullbacks). *Let $f(z) = e^z + c$ with $c \in J(f)$ non-recurrent. Then for any $z_0 \in \mathbb{C}$ there is a $\delta > 0$ such that for any $U \in \mathcal{C}_P(n, D_\delta(z_0))$, U is simply connected, and $f^n : U \rightarrow D_\delta(z_0)$ is univalent.*

Proof. Observe that for any bounded simply connected open set V not containing c , and any $U \in \mathcal{C}(1, V)$, U is simply connected, and $f : U \rightarrow V$ is univalent. In fact, $f : U \rightarrow V$ is a covering of a simply connected set, hence f is a universal covering and U is simply connected.

Fix $z_0 \in \mathbb{C}$. We first show that for any ε there exists δ , such that $\text{diam } U < \varepsilon$ for any $U \in \mathcal{C}(n, D_\delta(z_0))$ such that $c \in U$ and such that $f^n : U \rightarrow D_\delta(z_0)$ is univalent. If not there exists $\varepsilon > 0$, $\delta_k \rightarrow 0$, and $U_k \in \mathcal{C}(n_k, D_{\delta_k}(z_0))$ such that $\text{diam } U_k > \varepsilon$. By assumption the maps $\varphi_k : D_{\delta_k}(z_0) \rightarrow U_k$ are univalent, and for δ_k sufficiently small they all omit some specific periodic orbit of cardinality bigger than 3. The maps $\psi_k(z) = \varphi_k(\delta_k z + z_0) : \mathbb{D} \rightarrow U_k$ are normal by Montel Theorem, and they do not converge to infinity because $c \in U_k = \psi_k(\mathbb{D})$ for all k . So up to considering a subsequence they converge to a limit function ψ which is not constant because $\text{diam } U_k > \varepsilon$. As $c \in \psi_k(\mathbb{D})$ for all k , and ψ is not constant, there is a neighborhood V of c such that $V \subset \psi_k(\mathbb{D})$ for large k , hence $\psi_k^{-1}(V) \subset \mathbb{D}$. By definition of ψ_k , $\varphi_k^{-1}(V) = f^k(V) \subset D_{\delta_k}(z_0) \subset D_{\delta_0}(z_0)$ contradicting $c \in J(f)$.

Now let $D_\delta(z_0)$ be such that the previous claim holds for $\varepsilon = \text{dist}(c, \mathcal{P}(f))/2$, and let $U \in \mathcal{C}(n, D_\delta(z_0))$. If $c \notin f^j(U)$ for $j = 1 \dots n$, the claim follows by induction. Otherwise, whenever there exists j such that $c \in f^j(U)$, by the previous part of the proof $f^j(U) \cap \mathcal{P}(f) = \emptyset$, hence by forward invariance of $\mathcal{P}(f)$, $U \notin \mathcal{C}_P(n, D_\delta(x))$. \square

The next lemma is a tool to deal with sequences of pullbacks which go to infinity, i.e. do not have any finite accumulation point.

Lemma 3.2. *Let $f(z) = e^z + c$, $c \in J(f)$ non-recurrent. Let $z_0 \in \mathbb{C}$, $V = D_\delta(z_0)$ as in Lemma 3.1. Suppose that there exists a sequence $U_k \in \mathcal{C}_P(n_k, V)$, $k \rightarrow \infty$, with $\text{diam } U_k \geq \varepsilon_k > \delta/2^{n_k}$ for k sufficiently large. Then there exists a sequence of integers $j_k < n_k$ such that $U'_k \in \mathcal{C}_P(n_k - j_k - 1, V)$ has a finite accumulation point and*

$\text{diam } U'_k \geq \varepsilon'_k > e^{-M}(1 - e^{-\varepsilon_k})$. In particular, if the ε_k are bounded away from 0 so are the ε'_k , and if $\varepsilon_k \rightarrow \infty$, $\varepsilon'_k \rightarrow e^{-M}$. If $\inf \varepsilon_k > 0$, and $n_k \rightarrow \infty$, then $n_k - j_k \rightarrow \infty$.

Proof. Let $S = \{x, -M < \text{Re } x < \log 2\}$. Observe that $|f'(x)| \geq 2$ whenever $\text{Re } x \geq \log 2$, and that $\mathcal{P}(f) \cap U_k \neq \emptyset$ for all k , so, as $\text{diam } U_k > \delta/2^{n_k}$, there exists a minimal $j_k \geq 0$ such that $f^{j_k}(U_k) \cap S \neq \emptyset$.

Let $U'_k := f^{j_k+1}(U_k)$. As $f^{j_k}(U_k) \cap S \neq \emptyset$, the sets U'_k have a finite accumulation point in the annulus $A = f(S)$, centered at c , with internal radius e^{-M} and external radius 2. It is only left to estimate $\text{diam } U'_k$. As j_k is minimal, $\text{diam } f^{j_k}(U_k) \geq 2^{j_k} \text{diam } U_k \geq \varepsilon_k$. Between all sets of diameter L which intersect S and on which f is univalent, $|f'|$ is smallest along a horizontal segment of length L going to the left of S , whose image is a segment of length $e^{-M} - e^{-L-M}$, so $\text{diam } U'_k \geq e^{-M}(1 - e^{-\varepsilon_k})$.

To prove the last claim it is enough to show that the sequence $\{j_k\}$ is bounded. For each k , by minimality of j_k we have that $\text{diam } U_k \leq \text{diam } U'_k/2^{j_k}$; hence if $\inf \text{diam } U_k > 0$ and $j_k \rightarrow \infty$, $\text{diam } U'_k \rightarrow \infty$, contradicting normality of inverse branches on V together with the fact that $U'_k \cap A \neq \emptyset \forall k$. \square

Thanks to Lemma 3.2, the proofs of the following propositions and of Theorem A become similar to proofs from [ST] and [RvS]. We first need the following lemma ([ST, Lemma 2.1]).

Lemma 3.3. *For any $0 < \delta < 1$, there exists a constant $C(\delta)$ such that for any univalent map $g : U \rightarrow \mathbb{D}$ with U simply connected, and for any connected component U' of $g^{-1}(D_\delta(0))$, $\text{diam}_U(U') \leq C(\delta)$. Moreover $\lim_{\delta \rightarrow 0} C(\delta) = 0$.*

Proposition 3.4 (Uniform smallness). *Let $z_0 \in \mathbb{C}$, $D_\delta(z_0)$ be as in Proposition 3.1. Then for any $\varepsilon > 0$ there exists $D_{\delta'}(z_0) \subset D_\delta(z_0)$ such that for all $n \in \mathbb{N}$, $U \in \mathcal{C}_P(n, D_{\delta'}(z_0))$, $\text{diam } U < \varepsilon$.*

Proof. We first assume that there exists $R > 0$ such that $\text{diam } U < R$ for any $U \in \mathcal{C}_P(n, D_\delta(z_0))$. Then for any such U , $U \subset D_R(z)$ for some $z \in U$. On $D_R(z)$, the euclidean density is bounded by R times the hyperbolic density on $D_R(z)$. From this and monotonicity of the hyperbolic metric, we get that for any $U' \subset U \subset D_R(z)$

$$\text{diam } U' \leq R \text{diam}_{D_R(z)}(U') \leq R \text{diam}_U U' \leq \varepsilon$$

whenever δ' is sufficiently small so that $C(\delta') < \varepsilon/R$ as given by Lemma 3.3.

Let A be the annulus from Lemma 3.2. By normality of inverse branches defined on $D_\delta(z_0)$, there cannot be a sequence U_k intersecting A with $\text{diam } U_k \rightarrow \infty$, so there is some R such that $\text{diam } U < R$ for all U with $U \cap A \neq \emptyset$ and the claim holds for all U with $U \cap A \neq \emptyset$. It is only left to show that there is some R' such that $\text{diam } U < R'$ for all $U \in \mathcal{C}_P(n_k, D_{\delta'}(z))$, not necessarily intersecting A . Let δ' such that $\text{diam } U < e^{-M}/2$ for all $U \in \mathcal{C}_P(n, D_{\delta'}(z_0))$ intersecting A , and assume by contradiction that there exists

a sequence $U_k \in \mathcal{C}_P(n_k, D_{\delta'}(z))$ with $\text{diam } U_k \rightarrow \infty$. By Lemma 3.2, this gives a sequence of U'_k intersecting A with diameter tending to e^{-M} , contradicting the choice of δ' . The claim then holds up to making δ' smaller so that $C(\delta') < \min(\varepsilon/R', \varepsilon/R)$. \square

The proof of the next lemma is a combination of Lemma 3.2 with a classical lemma about backward shrinking (see [Ly1] for rational maps).

Proposition 3.5 (Backward shrinking). *Let $z \in J(f)$, $D_\delta(z)$ be as in Lemma 3.1. For all $\varepsilon > 0$ there exists n_ε , if $n > n_\varepsilon$ and $U \in \mathcal{C}_P(n, D_\delta(z))$, then $\text{diam } U < \varepsilon$.*

Proof. If not, there exist $\varepsilon > 0$ and a sequence $U_k \in \mathcal{C}_P(n_k, D_\delta(z))$, $n_k \rightarrow \infty$, with $\text{diam } U_k \geq \varepsilon$. As $n_k \rightarrow \infty$, eventually $\varepsilon/2^{n_k} < \delta$. Thanks to Lemma 3.2, up to replacing U_k with U'_k , we can assume that the U_k have a finite accumulation point y . By normality, the sequence of inverse branches $\varphi_k : D_\delta \rightarrow U_k$ has a limit function φ , which is non-constant because $\text{diam } U_k \geq \varepsilon$. It follows that there is a neighborhood of $\varphi(z)$ which is mapped inside $D_\delta(z)$ under infinitely many iterates of f , contradicting $\varphi(z) \cap J(f) \neq \emptyset$. \square

Proof of Theorem A. If $\mathcal{P}(f)$ is not hyperbolic, there are $n_k \rightarrow \infty$, and $z_k \in \mathcal{P}(f)$, such that $|(f^{n_k})'(z_k)| \leq 1$. As $\mathcal{P}(f)$ is closed and forward invariant, any $\{f^{n_k}(z_k)\}$, as well as any finite accumulation point thereof, belongs to $\mathcal{P}(f)$. Assume first that $f^{n_k}(z_k)$ has a finite accumulation point $y \in \mathcal{P}(f)$. By Propositions 3.1, 3.4 and 3.5, there is a neighborhood V of y such that for any $n \in \mathbb{N}$, $U \in \mathcal{C}_P(n, V)$

1. $\text{diam}(U) \leq \varepsilon$, and $f^n : U \rightarrow V$ is univalent;
2. $\forall \varepsilon' > 0 \exists n_0$, if $n \geq n_0$, $\text{diam}(U) \leq \varepsilon'$.

Up to considering a subsequence, we can assume that, for large k , $f^{n_k}(z_k) \in V$. For each such k , let U_k be the component of $f^{-n_k}(V)$ containing z_k . By local expansivity, $\text{diam}(f^j(U_k)) \leq \varepsilon$ for $j = 0 \dots n_k$, and $f^{n_k} : U_k \rightarrow V$ is univalent so it has a local inverse $\varphi_k : V \rightarrow U_k$. The family $\{\varphi_k\}$ is normal because it is a family of univalent inverse functions defined in V which omits at least three points; any limit function φ for φ_k is constant because by local expansivity $\text{diam}(U_k) \rightarrow 0$ as $k \rightarrow \infty$. This contradicts the initial assumption that

$$|\varphi'(y)| = \lim_{k \rightarrow \infty} |\varphi'_k(f^{n_k}(z_k))| = \lim_{k \rightarrow \infty} \left| \frac{1}{(f^{n_k})'(z_k)} \right| \geq 1.$$

Now suppose that $f^{n_k}(z_k) \rightarrow \infty$, let A, S be the annulus and the strip as in Lemma 3.2. To reduce to the previous case it is enough to find a sequence $j_k < n_k$ such that $j_k \rightarrow \infty$, $f^{j_k+1}(z_k) \in A$, and $|(f^{j_k+1})'(z_k)| \leq 1$.

If $f^{n_k}(z_k) \in A$, let $j_k = n_k$. Otherwise, for any k let $j_k \leq n_k - 1$ be the greatest integer such that $f^{j_k}(z_k) \in A$. Observe that the condition $|(f^{n_k})'(z_k)| \leq 1$, plus the fact that $f^{n_k}(z_k) \in \mathcal{P}(f)$, implies that such a j_k exists, otherwise we would have $|(f^{n_k})'(z_k)| \geq 2^{n_k} > 1$.

To prove that $|(f^{j_k+1})'(z_k)| \leq 1$, observe that

$$\begin{aligned} 1 &\geq |f^{n_k}(z_k)'| = \prod_{i=0}^{j_k} e^{\operatorname{Re} f^i(z_k)} \prod_{i=j_k+1}^{n_k-1} e^{\operatorname{Re} f^i(z_k)} \geq \\ &\geq \prod_{i=0}^{j_k} e^{\operatorname{Re} f^i(z_k)} 2^{n_k-j_k-1} \Rightarrow \prod_{i=0}^{j_k} e^{\operatorname{Re} f^i(z_k)} = |(f^{j_k+1})'(z_k)| \leq 1/2^{n_k-j_k-1} \leq 1. \end{aligned}$$

Similarly, because $\operatorname{Re} z > -M$ for all $z \in \mathcal{P}(f)$,

$$1 \geq |f^{n_k}(z_k)'| \geq \left(\prod_{i=0}^{j_k} e^{-M} \right) 2^{n_k-j_k-1} = e^{-M(j_k+1)} 2^{n_k-j_k-1}$$

hence, as $n_k \rightarrow \infty$, $j_k \rightarrow \infty$ as well. □

4 Puzzles and Parapuzzles

The construction of puzzles for polynomials (usually called Yoccoz Puzzle), as well as the construction of parapuzzles, has turned out to be very useful in proving rigidity results. Puzzles have been introduced in [BH], [Hu] and [Mi2]. Various versions have been constructed also for other maps, for example rational maps ([Ro]). In this section we describe the construction of puzzles and parapuzzles for the exponential family.

4.1 Puzzles and combinatorial non-recurrence

Let Γ be a forward invariant graph formed by finitely many periodic rays, and consider the connected components of $\mathbb{C} \setminus \Gamma$. For each n , the sets $\Gamma_n := f^{-n}(\Gamma)$ also partition \mathbb{C} into countably many components (see Figure 1 for an example of puzzle).

Definition 4.1. The countable collection of connected components of $\mathbb{C} \setminus f^{-n}(\Gamma)$ is called the *puzzle of level n* (induced by Γ); the connected components themselves are called *puzzle pieces of level n* and are denoted by $Y_j^{(n)}$. For each $n \in \mathbb{N}$ there is exactly one puzzle piece of level n containing $-\infty$, which is called the *branching puzzle piece* and denoted by Y_*^n . There is also exactly one *singular puzzle piece* containing c .

If c is non-escaping, it never belongs to Γ_n , hence at each level c belongs to the interior of some puzzle piece.

Remark 4.2. If c is a non-escaping parameter belonging to a parabolic wake attached to the period one hyperbolic component W_0 , there is a repelling fixed point $\alpha = \alpha(c)$, defined as an analytic continuation of the attracting fixed point, which is the landing point of finitely many periodic rays permuted transitively by the dynamics. These are the rays used for the most traditional puzzle construction. We use a more general Γ in order to allow for a weaker form of combinatorial non-recurrence (so that Theorem D holds). It is not known yet whether all non-escaping parameters are contained in such a wake, however, this is expected to be the case. A sufficient condition for this to happen is that $\mathcal{P}(f)$ is bounded (see [BL], Corollary 4.14).




Figure 1: Two possible topological configurations for the puzzle of level 4 with Γ formed by 3 rays of period 3 landing at a fixed point. The actual configuration depends on the position of c with respect to the puzzle of level 3. The configuration of puzzle pieces are invariant under translation by $2\pi i$. As the level increases, the new puzzle pieces are drawn in a paler color.

Remark 4.3. It is possible to define the label j for puzzle pieces of level n by using sequences in \mathbb{Z}^n , in a way that is consistent with the dynamics and with the vertical order of the dynamic rays at infinity. However, this takes some extra work and it is not needed for the rest of the paper; for simplicity, we will consider $j \in \mathbb{N} \cup \{*\}$.

The singular value c is called *combinatorially non-recurrent* if there exists some forward invariant graph Γ (formed by the closure of finitely many periodic dynamic rays) and some n such that the connected component of $\mathbb{C} \setminus \Gamma_n$ containing c does not intersect $\mathcal{P}(f)$.

Observe that parabolic and hyperbolic parameter are combinatorially recurrent. It is also known that the boundary of Siegel Disks is contained in the ω -limit set of a recurrent singular value (see [RvS], Corollary 2.10). So, in our setting, the condition of combinatorial non-recurrence implies that $J(f) = \mathbb{C}$. Combinatorial non-recurrence is a more restrictive condition than non-recurrence; however, we show in Section 7 that when $\mathcal{P}(f)$ is bounded and contained in the Julia set, non-recurrence implies combinatorial non-recurrence.

The next lemma follows directly from the definition of puzzle pieces, and the basic properties of dynamic rays.

Lemma 4.4. *The collections $\{Y_j^{(n)}\}$ define a puzzle for f , i.e.*

- $\forall n \in \mathbb{N}, i, j \in \mathbb{N} \cup \{*\}$, either $Y_j^{(n+1)} \subset Y_i^{(n)}$ or $Y_j^{(n+1)} \cap Y_i^{(n)} = \emptyset$.
- $\forall n, i \in \mathbb{N}$, $Y_i^{(n)}$ maps univalently to $Y_j^{(n-1)}$ for some j depending on i ;
- $Y_*^{(n)}$ is mapped to the singular puzzle piece of level $n - 1$ as an infinite degree covering.

Proof. Boundaries of puzzle pieces are dynamic rays, hence non-nested puzzle pieces do not intersect. Also, the boundaries of puzzle pieces of level $n + 1$ contain all preimages of the boundaries of puzzle pieces of level n , so the image of a puzzle piece cannot intersect two different puzzle pieces. \square

4.2 Parapuzzles and combinatorial equivalence

Let c_0 be a parameter, Γ be a forward invariant graph formed by periodic rays and their closure as in Section 4.1, and $\mathcal{A}_0 = \{\mathbf{s}_i\}_{i=1}^q$ be the set of addresses of the dynamic rays in Γ . Define

$$\mathcal{A}_k = \{\mathbf{s} \in \mathcal{S} : \sigma^j(\mathbf{s}) \in \mathcal{A}_0 \text{ for some } j \leq k\}.$$

The boundaries of puzzle pieces of level N consist exactly of dynamic rays with addresses in \mathcal{A}_N ; there rays are either periodic or preperiodic with preperiod $k \leq N$. Hence, by Propositions 2.5 and 2.6, the boundaries of puzzle pieces of level N move holomorphically in $\mathbb{C} \setminus \bigcup_{\mathbf{s} \in \mathcal{A}_N} \overline{G_{\mathbf{s}}}$.

A *parapuzzle piece* of level N for Γ is a region in parameter space over which the boundaries of puzzle pieces of level N move holomorphically.

Observe that the boundary of a parapuzzle piece of level N consists of countably many parameter rays with addresses in \mathcal{A}_N , hence of either parameter rays landing at Misiurewicz parameters or characteristic parameter rays. Also, parapuzzle pieces of different levels are either disjoint or contained one inside the other.

Two exponential maps $f_c, f_{c'}$ are *combinatorially equivalent* up to level N (with respect to the parapuzzle induced by Γ) if they belong to the same parapuzzle piece up to level N . If two maps are combinatorially equivalent as in Section 2.1, by Theorems 2.5 and 2.6 they belong to the same parapuzzle piece for all levels independently of the choice of Γ .

Two puzzle pieces $Y_\ell^{(n)}$ for f_c and $Y_\ell^{(n)'}$ for $f_{c'}$ are *equivalent* if they are bounded by dynamic rays with the same addresses which land together in the same pattern. If two maps are combinatorially equivalent up to level N , for each $n < N$ and for each puzzle piece $Y_\ell^{(n)}$ for f_c there is exactly one equivalent puzzle piece $Y_\ell^{(n)'}$ for $f_{c'}$. Equivalent puzzle pieces will be labelled with the same label.

Proposition 4.5 (Combinatorial equivalence). *If two exponential maps $f_c, f_{c'}$ are combinatorially equivalent up to level N , their singular values belong to equivalent puzzle pieces up to level $N - 1$.*

Proof. Suppose by contradiction that there is a level $n \leq N - 1$ such that c, c' belong to non-equivalent puzzle pieces. Then there are two addresses \mathbf{s}, \mathbf{s}' such that the curve $\overline{g_{\mathbf{s}}^c \cup g_{\mathbf{s}'}^c}$ encloses c , but the curve $\overline{g_{\mathbf{s}}^{c'} \cup g_{\mathbf{s}'}^{c'}}$ does not enclose c' . By Proposition 2.3, the puzzle pieces of level $n + 1$ which contain preimages of $g_{\mathbf{s}}^c, g_{\mathbf{s}'}^c$ on their boundary cannot be equivalent to the puzzle pieces of level $n + 1$ which contain preimages of $g_{\mathbf{s}}^{c'}, g_{\mathbf{s}'}^{c'}$; this contradicts combinatorial equivalence at level $n + 1$. \square

5 Quasiconformal Rigidity

In this section we prove Theorem B. We first construct a quasiconformal map ψ_N between f_c and $f_{c'}$, which is a conjugacy on the boundary of puzzle pieces up to a finite level N ; then ψ_N is used as the initial map in a lifting procedure to obtain a quasiconformal map Ψ which is a conjugacy on $\mathcal{P}(f)$; finally, Ψ is used in a new lifting procedure to obtain a quasiconformal map Φ which is a conjugacy on all preimages of $\mathcal{P}(f)$. In fact, because the latter are dense, by continuity Φ is a quasiconformal conjugacy on the entire plane.

Proposition 5.1 (Initial quasiconformal map). *Let c, c' be non-escaping and let $f_c, f_{c'}$ be combinatorially equivalent up to level N for some $N \in \mathbb{N}$. Then there exists a quasiconformal map $\psi_N : \mathbb{C} \rightarrow \mathbb{C}$ such that*

- $\psi_N(c) = c'$;
- ψ_N is a conjugacy on the dynamic rays $g_{\mathbf{s}}^c$ with $\mathbf{s} \in \mathcal{A}_k$ for all $k \leq N$.

Proof. As $f_c, f_{c'}$ are combinatorially equivalent up to level N , there is a parapuzzle piece Λ of level N which contains both c and c' . By Propositions 2.5 and 2.6 together

with Theorem 2.1, the map $h_{\tilde{c}} := g_{\mathbf{s}}^{\tilde{c}} \circ (g_{\mathbf{s}}^c)^{-1}$ defines a holomorphic motion over Λ of the dynamic rays $g_{\mathbf{s}}^c$ for $\mathbf{s} \in \mathcal{A}_j$ with $j \leq N$. By definition $h_{\tilde{c}}$ is a conjugacy on the dynamic rays $g_{\mathbf{s}}^c$ with $\mathbf{s} \in \mathcal{A}_j$ for all $j \leq N$. The singular value c itself also moves holomorphically in Λ , so $h_{\tilde{c}}$ can be defined at c as $h_{\tilde{c}}(c) = \tilde{c}$. By the Lambda Lemma (Theorem 1.6), $h_{\tilde{c}}$ can be extended to a quasiconformal map $\psi^{\tilde{c}} : \mathbb{C} \rightarrow \mathbb{C}$ for any $\tilde{c} \in \Lambda$. In particular, $\psi_N := \psi^{c'}$ is a conjugacy on the boundaries of puzzle pieces of level N and $\psi_N(c) = c'$. \square

From now on, let Γ be the forward invariant graph for which c is combinatorially non-recurrent, and consider the puzzle induced by Γ .

Proposition 5.2 (Conjugacy on $\mathcal{P}(f_c)$). *Let c, c' be non-escaping and let $f_c, f_{c'}$ be two combinatorially equivalent maps, such that c is combinatorially non-recurrent. Then there exists a quasiconformal map $\psi : \mathbb{C} \rightarrow \mathbb{C}$ which is a conjugacy on $\mathcal{P}(f_c)$.*

Proof. For each level n , denote by $P^{(n)}$ the union of the puzzle pieces of level n intersecting $\mathcal{P}(f_c)$. Observe that $P^{(n+1)} \subset P^{(n)}$, and that if a puzzle piece $Y \subset P^{(n+1)}$, then $f_c(Y) \subset P^{(n)}$. As f_c is combinatorially non-recurrent, there is some level N such that $c \notin P^{(N)}$. Let ψ_N be as in Proposition 5.1 and define inductively the sequence $\{\psi_n\}_{n \geq N}$ as:

$$\psi_{n+1} = \psi_n \quad \text{on } \mathbb{C} \setminus P^{(n)}; \quad (5.1)$$

$$\psi_{n+1} = f_{c'}^{-1} \circ \psi_n \circ f_c \quad \text{on } \overline{Y_\ell}, \text{ for any } Y_\ell = Y_\ell^{(n+1)} \subset P^{(n)}. \quad (5.2)$$

Observe that we are redefining ψ_n on all puzzle pieces of level $n+1$ which are contained in $P^{(n)}$, not only on the ones which are also contained in $P^{(n+1)}$. This is needed in order to ensure continuity. The branch of the inverse $f_{c'}^{-1}$ is defined so that $f_{c'}^{-1} \circ \psi_n \circ f_c(\overline{Y_\ell}) = \overline{Y_\ell}'$, where Y_ℓ' is the puzzle piece for $f_{c'}$ which is equivalent to Y_ℓ . We show by induction that this is possible while we prove the properties of $\{\psi_n\}$.

Now we show that the maps $\{\psi_n\}_{n \geq N}$ are well defined and satisfy the following properties:

1. $\psi_{n+1} = g_{\mathbf{s}}^{c'} \circ (g_{\mathbf{s}}^c)^{-1}$ on any $g_{\mathbf{s}}^c \in \partial Y^{(j+1)}$ such that $Y^{(j+1)} \subset P^{(j)}$ for some $j \leq n$.
2. ψ_{n+1} is K -quasiconformal.
3. $f_{c'} \circ \psi_{n+1} = \psi_n \circ f_c$ on $P^{(n+1)}$.

Observe that ψ_N satisfies all of the properties above for some quasiconformality constant K . We show by induction that they are satisfied for all $n > N$.

We first show that ψ_{n+1} is well defined on every $Y_\ell^{(n+1)} \subset P^{(n)}$. The image $f_c(Y_\ell^{(n+1)}) = \overline{Y_\ell^{(n)}}$ for some puzzle piece $Y_\ell^{(n)}$ of level n . Let $Y_\ell'^{(n)}$ be the puzzle

piece which is equivalent to $Y_{\bar{\ell}}^{(n)}$. The puzzle piece $\overline{Y_{\bar{\ell}}^{(n)}}$ does not contain c by combinatorial non-recurrence, so $\overline{Y_{\bar{\ell}}^{(n)}}$ does not contain c' by Proposition 4.5. In particular inverse branches for $f_{c'}$ are well defined and univalent in a neighborhood of $\overline{Y_{\bar{\ell}}^{(n)}}$, and there is a univalent branch $f_{c'}^{-1} : Y_{\bar{\ell}}^{(n)} \rightarrow Y_{\bar{\ell}}^{(n+1)}$.

We start by showing Property 1. Let $g_{\bar{s}}^c$ be a dynamic ray such that $g_{\bar{s}}^c \in \partial Y^{(j+1)}$ for some $Y^{(j+1)} \subset P^{(j)}$ and some $j \leq n$. Then $g_{\sigma\bar{s}}^c \in \partial Y^{(j)}$ for some $Y^{(j)} \subset P^{(j-1)}$. By direct computation and by the induction hypothesis, on $g_{\bar{s}}^c$,

$$\psi_{n+1}|_{g_{\bar{s}}^c} = f_{c'}^{-1} \circ \psi_n \circ f_c|_{g_{\bar{s}}^c} = \tag{5.3}$$

$$= f_{c'}^{-1} \circ g_{\sigma\bar{s}}^{c'} \circ (g_{\sigma\bar{s}}^c)^{-1} \circ f_c|_{g_{\bar{s}}^c} \tag{5.4}$$

$$= g_{\bar{s}}^{c'} \circ (g_{\bar{s}}^c)^{-1}|_{g_{\bar{s}}^c}. \tag{5.5}$$

(The second equality follows from Property 1. in the induction hypothesis, while the last equality comes from the functional equation in Theorem 2.1 and the choice of branch of $f_{c'}^{-1}$). In particular, by the induction hypothesis $\psi_{n+1} = \psi_n$ on any $g_{\bar{s}}^c \in \partial Y^{(j+1)}$ if $Y^{(j+1)} \subset P^{(j)}$ for $j \leq n-1$.

Property 1. implies continuity: on rays belonging to $\partial P^{(n)}$ we obtain that $\psi_{n+1} = \psi_n$, while on any ray $g_{\bar{s}}^c$ in the boundary of two puzzle pieces of level $n+1$ the new defined functions match because ψ_{n+1} coincides with $g_{\bar{s}}^{c'} \circ (g_{\bar{s}}^c)^{-1}$. So ψ_{n+1} is a homeomorphism; to show that it is K -quasiconformal, note that whenever ψ_{n+1} is redefined, it is done by pre-post composing with a conformal map, so the dilatation remains K and ψ_{n+1} is K -quasiconformal (as it is a homeomorphism which is K -quasiconformal outside a countable number of smooth curves). This proves Property 2.. Property 3. follows directly from the definition of ψ_{n+1} because $P^{(n+1)} \subset P^{(n)}$.

By Property 2. the functions ψ_n are uniformly quasiconformal, can be extended as to fix infinity and by Property 3. they all coincide on $\partial P^{(N)}$, so by Lemma 1.2 there exists a K -quasiconformal limit function $\psi : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$ fixing infinity. By hyperbolicity of $\mathcal{P}(f)$ the limit is unique. By Property 3., the limit map ψ is a conjugacy between f_c and $f_{c'}$ on $\bigcap_{n \geq N} P^{(n)} = \mathcal{P}(f)$. \square

We conclude the proof of Theorem B by lifting ψ to a continuous map which is a conjugacy on the preimages of the postsingular set, and then use the fact that the latter are dense to obtain by continuity a conjugacy on the entire plane. The next theorem is one of the fundamental facts in the theory of covering spaces (see e.g. [Ha], Proposition 1.33).

Theorem 5.3 (Lifting). *Let X, Y, Z be topological spaces with base points x, y, z respectively. If X is a covering of Z via a map h such that $h(x) = z$, and $f : Y \rightarrow Z$ is a continuous map with $f(y) = z$ and Y is simply connected, then there exists a unique continuous lift $\tilde{f} : Y \rightarrow X$ with $\tilde{f}(y) = x$.*

Proof of Theorem B. Let Ψ_0 be ψ as given by Proposition 5.2. Consider the lifting diagram below (where the couples (X, x) mean the space X with base point x):

$$\begin{array}{ccc} & \Psi_{n+1} & \\ (\mathbb{C}, c) & \longrightarrow & (\mathbb{C}, c') \\ f_c \downarrow & & \downarrow f_{c'} \\ (\mathbb{C} \setminus \{c\}, f_c(c)) & \xrightarrow{\Psi_n} & (\mathbb{C} \setminus \{c'\}, f_{c'}(c')) \end{array}$$

As \mathbb{C} is simply connected, the existence of the maps Ψ_n as lift of $f_c \circ \psi_n$ is ensured by Theorem 5.3, and for each n , $\Psi_n(c) = c'$. The existence of Ψ_{n+1} gives, for any $z \in \mathbb{C}$, a preferred choice of an inverse branch for f_c^{-1} in the definition of $\Psi_{n+1}(z) = (f_{c'}^{-1} \circ \Psi_n \circ f_c)(z)$.

Let us show by induction that for each n , Ψ_n is a conjugacy on $\bigcup_{j \leq n} f_c^{-j}(\mathcal{P}(f_c))$. This is verified for $n = 0$ by Proposition 5.2. Assume by induction that Ψ_n is a conjugacy on $\bigcup_{j \leq n} f_c^{-j}(\mathcal{P}(f_c))$. We first show that for $z \in f_c^{-j}(\mathcal{P}(f_c))$, $j \leq n$, $\Psi_{n+1}(z) = \Psi_n(z)$. In fact,

$$\Psi_{n+1}(z) = (f_{c'}^{-1} \circ \Psi_n \circ f_c)(z) = (f_{c'}^{-1} \circ f_{c'} \circ \Psi_n)(z) = \Psi_n(z). \quad (5.6)$$

(The last equality holds by choice of $f_{c'}^{-1}$). In particular, Ψ_{n+1} is still a conjugacy on $\bigcup_{j \leq n} f_c^{-j}(\mathcal{P}(f_c))$. To show that Ψ_{n+1} is a conjugacy also on $f_c^{-n-1}(\mathcal{P}(f_c))$, let $z \in f_c^{-n-1}(\mathcal{P}(f_c))$; by definition

$$f_{c'} \circ \Psi_{n+1} = \Psi_n \circ f_c = \Psi_{n+1} \circ f_c$$

by Equation 5.6, because $f_c(z) \in f_c^{-n}(\mathcal{P}(f_c))$.

As locally Ψ_{n+1} is obtained by pre- and post-composing Ψ_n with conformal maps, all Ψ_n are uniformly quasiconformal, and by Equation 5.6 they all coincide on the postsingular set. Extending the Ψ_n to fix infinity, Lemma 1.2 gives a limit map Ψ which is a conjugacy on $\bigcup_{n \in \mathbb{N}} f_c^{-n}(\mathcal{P}(f_c))$. The latter ones are dense, so by continuity Ψ is a conjugacy on all of \mathbb{C} . \square

6 Conformal Rigidity

When the postsingular set is bounded, the proof of Theorem C can be made relatively easy by using Theorem 1.4.

Proof of Theorem C for bounded postsingular set. Let Ψ be the quasiconformal conjugacy obtained in Theorem B, and σ_0 be the standard conformal structure on \mathbb{C} . The pushforward of σ_0 by Ψ defines an invariant conformal structure σ' in the dynamical plane for $f_{c'}$. The conformal structure σ' defines an invariant line field, which is constant by Theorem 1.4. So σ' is the standard conformal structure and Ψ is conformal by

Weyl's Lemma, hence $c' = c + 2\pi in$ for some n . As c, c' are combinatorially equivalent, they are in the same fiber, and $c = c'$. \square

A more involved proof, using an open-closed argument, shows that the reduced fibers of combinatorially non-recurrent parameters are trivial also when the postsingular set is not bounded. Let $\text{QC}(c)$ be the quasiconformal class of c , that is the connected component containing c of the set of parameters c' such that $f_{c'}$ is quasiconformally conjugate to f_c . Observe that parameters of the form $c + 2\pi in$ cannot belong to $\text{QC}(c)$ unless $n = 0$.

Lemma 6.1 (Quasiconformal classes are open). *Let c be non-escaping and combinatorially non-recurrent. If $\text{QC}(c) \neq \{c\}$, $\text{QC}(c)$ is open.*

Proof. Let $c \neq c' \in \text{QC}(c)$, Ψ be a quasiconformal conjugacy between f_c and $f_{c'}$. Let $\mu_0 = 0$ be the Beltrami coefficient for the standard conformal structure, $\mu' := \Psi_*\mu_0$ the Beltrami coefficient obtained in the dynamical plane for $f_{c'}$, and for $\lambda \in \mathbb{D}$ let μ_λ over $\lambda \in \mathbb{D}$ be an analytic interpolation between μ and μ' (for example, $\mu_\lambda = \lambda\mu'$). If $\mu' \neq 0$, hence the deformation is non-trivial, by the Measurable Riemann Mapping Theorem to each μ_λ corresponds a quasiconformal map Ψ_λ conjugating f_c to an exponential map f_λ . As μ_λ depends holomorphically on λ , the Ψ_λ , and hence f_λ , depend holomorphically on λ ; in particular there is an open neighborhood of c on which all maps are quasiconformally conjugate to f_c . With the same argument, it is possible to find an open neighborhood contained in $\text{QC}(c)$ for any $\tilde{c} \in \text{QC}(c)$. \square

A maximal open set of parameters all of which are topologically conjugate to each other, and which are not hyperbolic, is called a *non-hyperbolic component*. Any such component \mathcal{Q} is simply connected; otherwise, there would be bounded components of $\mathbb{C} \setminus \overline{\mathcal{Q}}$, contradicting the fact that escaping parameters are dense in the bifurcation locus.

Lemma 6.2 (Boundaries of non-hyperbolic components). *Consider the exponential family. The boundary of a non-hyperbolic component \mathcal{Q} in parameter space cannot contain escaping parameters which are accessible from the inside of \mathcal{Q} . In particular, $\partial\mathcal{Q}$ contains infinitely many non-escaping parameters on its boundary.*

Proof. As \mathcal{Q} is simply connected it can be uniformized to the unit disk, and by the boundary behavior of the Riemann map, there is only a set of measure zero of parameters which are not accessible from the inside of \mathcal{Q} . Also, non-hyperbolic components are fully contained in a single fiber because they cannot intersect parameter rays, hence by Lemma 2.7, $\overline{\mathcal{Q}}$ intersects only finitely many parameter rays. So there are infinitely many parameters on $\partial\mathcal{Q}$ which are accessible via a curve $\gamma : [0, 1) \rightarrow \mathcal{Q}$, and which are not endpoints of parameter rays. To prove the claim it is enough to show that no such accessible parameter can be escaping. Suppose by contradiction that $c_0 \in \partial\mathcal{Q}$ is an escaping parameter accessible from the inside of \mathcal{Q} , and is not the endpoint of

a parameter ray. Then $c_0 = G_{s_0}(t_0)$ for some parameter ray G_{s_0} , and there exists an arc $G_{s_0}(t_0 - \varepsilon, t_0 + \varepsilon)$ which can be oriented following increasing t . Also, G_{s_0} is approximated on compact sets on both sides by curves in hyperbolic components (see Section 2.2); as γ is a local transversal to the arc $G_{s_0}(t_0 - \varepsilon, t_0 + \varepsilon)$ at c_0 , infinitely many of these curves have to intersect γ , which is impossible because \mathcal{Q} is a non-hyperbolic component. \square

Proof of Theorem C. Let c, c' be non-escaping and combinatorially equivalent, c be combinatorially non-recurrent. Then $c' \in \text{QC}(c)$ by Theorem B, and $c \in \mathcal{Q}(c')$ for some non-hyperbolic component $\mathcal{Q}(c')$, because quasi-conformal conjugacy implies topological conjugacy, and because combinatorial non-recurrence implies that c is not hyperbolic. For the same reason, $\text{QC}(c) \subset \mathcal{Q}(c)$. Let $F_R(c)$ denote the reduced fiber of c . As parameter rays cannot intersect non-hyperbolic components, $\mathcal{Q}(c) \subset F_R(c)$. On the other side, by Theorem B, $F_R(c) \subset \text{QC}(c)$, hence $\text{QC}(c) = \mathcal{Q}(c) = F_R(c)$. $F_R(c)$ is open by Lemma 6.1, while it contains at least one of its boundary points by Lemma 6.2, giving a contradiction. \square

If two parameters are in the same fiber, then they are combinatorially equivalent, so Theorem C implies that reduced fibers of combinatorially non-recurrent parameters are trivial.

7 Non-recurrence and combinatorial non-recurrence

In this section we show that for parameters with bounded postsingular set, non-recurrence implies combinatorial non-recurrence provided they are not hyperbolic or parabolic parameters. We achieve this through the combinatorial similarity between the parameter plane for exponentials and for unicritical polynomials. We use several results about polynomial dynamics which are most likely known to the reader, like the analogues of the theorems described for exponentials in Section 4 (see e.g. [Mi]). We add references for the less well known results. To introduce notation, we recall the definitions of dynamic and parameter rays for polynomials.

Let $P_c^D(z) = z^D + c$ be a unicritical polynomial of degree D . If c is non-escaping, the complement of the filled Julia set K_c can be uniformized by $\mathbb{C} \setminus \overline{\mathbb{D}}$ via the Böttcher map, which is also a conjugacy between P_c^D and P_0^D in a neighborhood of infinity. The angles of the straight rays in $\mathbb{C} \setminus \overline{\mathbb{D}}$ can be expressed as sequences \mathbf{s} over an alphabet of D symbols, containing integer entries $s_i \in (-D/2 + 1/2, +D/2 + 1/2)$ if D is even, and $s_i \in (-D/2, +D/2)$ if D is odd. The preimage of a straight ray of angle \mathbf{s} under the Böttcher map is called the *dynamic ray* of angle \mathbf{s} and denoted by $g_{\mathbf{s}}^{D,c}$ or just $g_{\mathbf{s}}^c$ when the degree is implicit. The dynamics of P_c^D on the dynamic rays is conjugate to the dynamics of the shift map $\sigma = \sigma_D$ over the sequences over D symbols. Similarly, the complement of the connectedness locus M_D for the family $\{P_c^D\}$ can be uniformized

by $\mathbb{C} \setminus \overline{\mathbb{D}}$, and the preimage of a straight ray of angle \mathbf{s} is called the *parameter ray* $G_{\mathbf{s}}^D$ of angle \mathbf{s} . Near infinity, both dynamic and parameter rays respect the *cyclic order* induced by the cyclic order on the set of sequences over D symbols identified with the unit circle.

7.1 Proof of Theorem D

For the proof of Theorem D we need some additional results about the exponential family as well as some rather specific knowledge about the parameter structure of unicritical polynomials. We tried to make this section as self-contained as possible, avoiding however to dwell excessively in the theory of renormalization for unicritical polynomials.

We make use of the following two results from [BL] (see [BL, Corollary 4.6] and [BL, Corollary 4.14] respectively).

Theorem 7.1 (Accessibility of c). *If f_c is an exponential map with $c \in J(f_c)$ and $\mathcal{P}(f_c)$ is bounded, then there is at least a dynamic ray $g_{\mathbf{s}}$ landing at c , and $\|\mathbf{s}\| < \infty$. Also, the length of the arcs $g_{\sigma^n \mathbf{s}}(0, t) \rightarrow 0$ uniformly in n as $t \rightarrow 0$.*

The last estimate and the fact that $\|\mathbf{s}\| < \infty$ are not stated explicitly, but they follow directly from the construction.

Theorem 7.1 is the only place in the proof of Theorem D where boundedness of $\mathcal{P}_c(f)$ is used, and are expected to hold in the general non-recurrent case.

An address or angle \mathbf{s} is called *non-recurrent* if $\mathbf{s} \notin \overline{O(\mathbf{s})} := \overline{\{\sigma^n(\mathbf{s})\}}_{n \in \mathbb{N}}$.

Lemma 7.2. *Let $f_c(z) = e^z + c$ or $f_c(z) = z^D + c$ be non-recurrent. If there is a dynamic ray $g_{\mathbf{s}}$ landing at c such that $g_{\sigma^n \mathbf{s}}(0, t) \rightarrow 0$ uniformly in n as $t \rightarrow 0$, then \mathbf{s} is non-recurrent.*

Proof. By non-recurrence there is a disk $D_\varepsilon(c)$ such that $D_\varepsilon(c) \cap \mathcal{P}(f_c) = \emptyset$. As $g_{\mathbf{s}}$ lands at c , for any $k \in \mathbb{N}$ the dynamic ray $g_{\sigma^k \mathbf{s}}$ lands at $f_c^k(c)$. Let t_ε be such that

$$\ell(g_{\sigma^n \mathbf{s}}(0, t)) < \varepsilon/2 \tag{7.1}$$

for all $n \in \mathbb{N}, t < t_\varepsilon$. Also let $t_0 < t_\varepsilon$ such that $g_{\mathbf{s}}(t_0) \in D_{\varepsilon/2}(c)$.

If \mathbf{s} is recurrent, there is a subsequence $\sigma^k \mathbf{s} \rightarrow \mathbf{s}$ as $k \rightarrow \infty$, hence by Lemma 2.2, $g_{\sigma^k \mathbf{s}}(t_0) \rightarrow g_{\mathbf{s}}(t_0)$ as $k \rightarrow \infty$. By Equation 7.1, $g_{\sigma^k \mathbf{s}}$ eventually lands inside $D_\varepsilon(c)$, contradicting $D_\varepsilon(c) \cap \mathcal{P}(f_c) = \emptyset$. \square

Lemma 7.3. *Let f_c be a unicritical polynomial or an exponential map. If a parameter ray $G_{\mathbf{s}}$ lands at a parameter c and $c \in J(f_c)$, then $g_{\mathbf{s}}^c$ belongs to the dynamic fiber of c . Viceversa, if $g_{\mathbf{s}}^c$ lands at c , the parameter ray $G_{\mathbf{s}}$ belongs to the parameter fiber of c .*

Proof. Suppose that $G_{\mathbf{s}}$ lands at c and that $g_{\mathbf{s}}^c$ does not belong to the dynamic fiber of c . Then there is a pair of dynamic rays $g_{\mathbf{s}^+}^c, g_{\mathbf{s}^-}^c$ separating $g_{\mathbf{s}}^c$ from c , whose addresses are either periodic or preperiodic. If the addresses are periodic, there are also two characteristic rays enclosing c which separate $g_{\mathbf{s}}^c$ from c , and the parameter rays $G_{\mathbf{s}^+}, G_{\mathbf{s}^-}$ form a parabolic wake in parameter plane Proposition 2.5. If the addresses are preperiodic, the parameter rays $G_{\mathbf{s}^+}, G_{\mathbf{s}^-}$ form a Misiurewicz wake by Proposition 2.6. In both cases, the wake separates $G_{\mathbf{s}}$ from c by vertical order of parameter rays in parameter plane, so $G_{\mathbf{s}}$ does not belong to the fiber of c . The case in which $g_{\mathbf{s}}^c$ lands at c and $G_{\mathbf{s}}$ does not belong to the fiber of c is analogous. \square

We now spend some time proving that for unicritical polynomials parameter rays with non-recurrent addresses land at non-recurrent parameters.

Proposition 7.4 (Landing of non-recurrent rays). *Let $G_{\mathbf{s}}^D$ be a parameter ray of angle \mathbf{s} in the parameter plane for the family $\{P_c^D\}$. If \mathbf{s} is non-recurrent, then $G_{\mathbf{s}}^D$ lands at a parameter \tilde{c} which is at most finitely renormalizable and non-recurrent.*

What follows is a brief introduction to renormalization and rigidity for unicritical polynomials. A unicritical polynomial $P_c^D(z) = z^D + c$ with connected Julia set is called *renormalizable of period n* if there are neighborhoods U, U' of 0, with U compactly contained in U' , such that $f^n : U \rightarrow U'$ is a degree D polynomial-like map with connected Julia set (see e.g. [Hu]). The new polynomial-like map can be itself renormalizable, and so on.

For quadratic polynomials, the connected components of sets of parameters which are renormalizable of period n form small copies of the Mandelbrot set (see [DH]). Parameters which are renormalizable of period n , and hence these copies, are combinatorially characterized by being contained in some parabolic wake bounded by two characteristic angles $\mathbf{s}^+, \mathbf{s}^-$, from which infinitely many Misiurewicz wakes have been cut out. We will call this the *renormalization wake* of the corresponding small copy. The dynamic rays which are left in this renormalization wake are exactly the ones whose angles are represented by sequences in $\{t^+, t^-\}^{\mathbb{N}}$, where t^+, t^- are the finite sequences of n symbols with $\mathbf{s}^+ = \overline{t^+}$, $\mathbf{s}^- = \overline{t^-}$. This is the same as saying that the angle is the output of a *tuning* through the angles $\mathbf{s}^+, \mathbf{s}^-$ as described in [Do]. Observe that a parameter ray accumulates on a given small copy if and only if it belongs to its renormalization wake.

For unicritical polynomials of higher degree D , renormalizable parameters of period n also form small copies of the connectedness locus M_D and can be characterized in a similar way (see [Sc1, Theorem 3.1]). The renormalization wake is again formed by a parabolic wake, from which countably many Misiurewicz wakes have been cut out. The angles of the parameter rays contained in the renormalization wake can be describes as the sequences in $\{t^+, t^-, t^3 \dots t^{D-2}\}^{\mathbb{N}}$ where t^i are specific finite sequences of length n . Again, a parameter ray accumulates on a given small copy if and only if it belongs to its renormalization wake.

An angle is called *combinatorially renormalizable* of period n if it belongs to the renormalization wake of a small copy of renormalization period n .

Lemma 7.5. *If \mathbf{s} is a non-recurrent angle written in D -adic expansion, then it is at most finitely many times combinatorially renormalizable (when seen as the angle of a parameter ray for polynomials of degree D).*

Proof. If \mathbf{s} is renormalizable of some period q , then \mathbf{s} is represented by a sequence constructed out of only D blocks of length q , hence there are infinitely many k such that $|\mathbf{s} - \sigma^k \mathbf{s}| < \frac{2}{D^q}$. If \mathbf{s} is infinitely many times combinatorially renormalizable, then there exists a sequence $q_n \rightarrow \infty$ and $k_n \rightarrow \infty$ such that $|\mathbf{s} - \sigma^{k_n} \mathbf{s}| < \frac{2}{D^{q_n}}$, contradicting non-recurrence of \mathbf{s} . \square

The next theorem is one of the cornerstones in the theory of rigidity for quadratic and unicritical polynomials (see [Hu], [AKLS] respectively).

Theorem 7.6 (Yoccoz Theorem). *Let P_c^D be an at most finitely renormalizable unicritical polynomial with all periodic points repelling and connected Julia set. Then P_c^D is combinatorially rigid, or equivalently the reduced fiber of c is a single point.*

The condition of combinatorial rigidity is often stated as *parapuzzle pieces shrink to points*. To prove Yoccoz Theorem local connectivity of the Julia set is needed. For unicritical polynomials see [KL, Theorem A]:

Theorem 7.7. *Let P_c^D be an at most finitely renormalizable unicritical polynomial with all periodic points repelling and connected Julia set. Then $J(P_c^D)$ is locally connected.*

By Caratheodory's Theorem, local connectivity of the Julia set implies that all dynamic rays land and all points are landing points of a dynamic ray. In particular, the critical value is accessible. It is also known that for a non-renormalizable unicritical polynomial with all periodic points repelling and connected Julia set, the puzzle pieces induced by the periodic rays landing at the α -fixed points shrink to points. So, for an at most finitely renormalizable polynomial of the same kind, the puzzle pieces induced by some forward invariant graph $\tilde{\Gamma}$ formed by finitely many periodic rays also shrink to points.

After the preparation about tuning, Proposition 7.4 is essentially a consequence of Yoccoz's Theorem.

Proof of Proposition 7.4. By Lemma 7.5, \mathbf{s} is only finitely renormalizable, so $G_{\mathbf{s}}$ belongs to at most finitely many renormalization wakes and can accumulate only on non-escaping parameters which are at most finitely renormalizable. By Yoccoz Theorem, the reduced fiber of any such parameter \tilde{c} is trivial, hence if $G_{\mathbf{s}}^D$ accumulates on \tilde{c} in fact it lands at it. By Proposition 7.3, $g_{\mathbf{s}}^{\tilde{c}}$ belongs to the dynamical fiber of \tilde{c} , and by local connectivity of $J(f_{\tilde{c}})$ in fact it lands at \tilde{c} . As \mathbf{s} is a non-recurrent address, \tilde{c} is non-recurrent. \square

We present two more results and then prove Theorem D.

Proposition 7.8 ([Be, Theorem 4.7]). *Let $\mathbf{s}_1 \dots \mathbf{s}_q$ be a finite set of preperiodic addresses. Then the parameter rays of addresses $\mathbf{s}_1 \dots \mathbf{s}_q$ land together at a Misiurewicz parameter in the exponential parameter plane if and only if they land together at a Misiurewicz parameter in the parameter plane of any family of unicritical polynomials with sufficiently high degree D .*

The next proposition is a direct consequence of [Be, Theorem 4.11] and Theorem 2.5.

Proposition 7.9. *Two parameter rays with periodic addresses $\mathbf{s}^+, \mathbf{s}^-$ land together in the parameter plane for exponentials if and only if the parameter rays with angles $\mathbf{s}^+, \mathbf{s}^-$ land together in the parameter plane for all families $\{P_c^D\}$ with sufficiently high degree D .*

Proof of Theorem D. Recall that f_c is an exponential map such that $\mathcal{P}(f_c) \subset J(f_c)$ is bounded. Let $g_{\mathbf{s}}$ be the dynamic ray landing at c given by Theorem 7.1 and let D be sufficiently large. By Lemma 7.2, \mathbf{s} is non-recurrent, so by Proposition 7.4 the parameter ray $G_{\mathbf{s}}^D$ (for the family of unicritical polynomials of degree D) lands at a finitely renormalizable non-recurrent polynomial parameter \tilde{c} .

By Lemma 7.3, the dynamic ray $g_{\mathbf{s}}^{\tilde{c}}$ belongs to the dynamical fiber of \tilde{c} . As \tilde{c} is at most finitely renormalizable, by Theorem 7.7 its Julia set is locally connected, so in fact $g_{\mathbf{s}}^{\tilde{c}}$ lands at \tilde{c} . Because \tilde{c} is at most finitely renormalizable, there is some n and some cycle $\tilde{\Gamma}$ of rays of period n such that the puzzle pieces induced by $\tilde{\Gamma}$ and containing c shrink to points.

By non-recurrence of \tilde{c} , $\text{dist}(\tilde{c}, \mathcal{P}(f_{\tilde{c}})) > 0$, hence there is some level of the puzzle induced by $\tilde{\Gamma}$ for which the singular value and $\mathcal{P}(f_{\tilde{c}})$ are separated. This means that there are finitely many preperiodic rays $\{g_{\mathbf{s}_i}^{\tilde{c}}\}_{\mathbf{s}_i \in \mathcal{K}}$ separating \tilde{c} from $\mathcal{P}(f_{\tilde{c}})$.

By the polynomial analogue of Propositions 2.5 and 2.6, \tilde{c} is contained in finitely many wakes defined by the parameter rays $\{G_{\mathbf{s}_i}^D\}_{\mathbf{s}_i \in \mathcal{K}}$ (it can be contained in many more wakes, but this is irrelevant to us).

By Propositions 7.8 and 7.9, the same wakes exist in the exponential parameter plane, and by vertical order of parameter rays, the parameter ray $G_{\mathbf{s}}$ (and hence c by Lemma 7.3) is contained in all of them. By Proposition 2.5 and 2.6, the corresponding pairs of rays with addresses in \mathcal{K} land together in the dynamical plane for c . By vertical order of dynamical rays, as the rays $g_{\mathbf{s}_i}^{\tilde{c}}$ separate \tilde{c} from its forward orbit in the polynomial dynamical plane (and each point on the orbit is the landing point of a dynamic ray $g_{\sigma^k \mathbf{s}}^{\tilde{c}}$ by Theorem 7.1), the rays $\{g_{\mathbf{s}_i}^c\}_{\mathbf{s}_i \in \mathcal{K}}$ also separate c from its forward orbit in the exponential plane. By continuity, they also separate c from the closure of its forward orbit i.e. from the postsingular set. \square

References

- [Ah] L. V. Ahlfors, *Lectures on quasiconformal mappings*. University Lecture Series, 38. American Mathematical Society, Providence, RI, (2006).
- [AKLS] A. Avila, J. Kahn, M. Lyubich, W. Shen, *Combinatorial rigidity for unicritical polynomials*. Ann. of Math. (2) 170 (2009), no. 2, 783-797.
- [BR] N. I. Baker, P. J. Rippon, *Iteration of exponential functions*. Ann. Acad. Sci. Fenn. Ser. A I Math. 9 (1984), 49–77.
- [Be] A. M. Benini, *Triviality of fibers for Misiurewicz parameters in the exponential family*. Conform. Geom. Dyn. 15 (2011), 133-151 .
- [BL] A. M. Benini, M. Lyubich, *Repelling periodic points and landing of rays for post-singularly bounded exponential maps*. arXiv:1208.0147 [math.DS].
- [BH] B. Branner, J. H. Hubbard, *The iteration of cubic polynomials, Part II*. Acta Math. v. 169 (1992), 229–325.
- [BDHRGH1] C. Bodelón, R. L. Devaney, M. Hayes, G. Roberts, L. R. Goldberg, J. H. Hubbard, *Hairs for the complex exponential family*. Internat. J. Bifur. Chaos Appl. Sci. Engrg. 9 (1999), no. 8, 1517–1534.
- [BDHRGH2] C. Bodelón, R. L. Devaney, M. Hayes, G. Roberts, L. R. Goldberg, J. H. Hubbard, *Dynamical convergence of polynomials to the exponential*. J. Differ. Equations Appl. 6 (2000), no. 3, 275–307.
- [Do] A. Douady, *Algorithms for computing angles in the Mandelbrot set*. Chaotic dynamics and fractals (Atlanta, 1985), 155-168.
- [DH] A. Douady, J. H. Hubbard, *On the dynamics of polynomial-like mappings*. Ann. Sci. École Norm. Sup. (4) 18 (1985), no. 2, 287-343.
- [EL] A. Eremenko, M. Lyubich, *Dynamical properties of some classes of entire functions*. Ann. Inst. Fourier (Grenoble) 42 (1992), no. 4, 989–1020.
- [FS] M. Förster, D. Schleicher, *Parameter rays in the space of exponential maps*. Ergodic Theory Dynam. Systems 29 (2009), no. 2, 515–544.
- [GKS] J. Graczyk, J. Kotus, G. Świątek, *Non-recurrent meromorphic functions*. Fund. Math. 182 (2004), no. 3, 269–281.
- [Ha] A. Hatcher, *Algebraic topology*. Cambridge University Press, Cambridge, 2002.
- [Hu] J. H. Hubbard, *Local connectivity of Julia sets and bifurcation loci: three theorems of J. C. Yoccoz*. Topological Methods in Modern Mathematics, ed. Goldberg and Phillips, Publish or Perish (1993), 467-511.
- [KL] J. Kahn, M. Lyubich, *Local connectivity of Julia sets for unicritical polynomials*. Ann. of Math. (2) 170 (2009), no. 1, 413-426.

- [LSV] B. Laubner, D. Schleicher, V. Vicol, *A combinatorial classification of postsingularly finite complex exponential maps*. Discrete Contin. Dyn. Syst. 22 (2008), no. 3, 663–682.
- [Ly1] M. Lyubich, *Entropy properties of rational endomorphisms of the Riemann sphere*. Ergodic Theory Dynam. Systems 3 (1983), no. 2, 351–385.
- [Ly2] M. Lyubich, *Dynamics of quadratic polynomials I-II*. Acta Math. 178 (1997), 185–297.
- [MS] P. Makienko, G. Sienna, *Poincaré series and instability of exponential maps*. Bol. Soc. Mat. Mexicana (3) 12 (2006), no. 2, 213–228.
- [Ma] R. Mañé, *On a theorem of Fatou*. Bol. Soc. Brasil. Mat. (N.S.) 24 (1993), no. 1, 1–11.
- [MSS] R. Mañé, P. Sad, and D. Sullivan, *On the dynamics of rational maps*. Ann. Sci. École Norm. Sup. (4) 16 (1983), no. 2, 193–217.
- [Mi] J. Milnor, *Dynamics in one complex variable*. Annals of Mathematics Studies (2006), Princeton University Press.
- [Mi2] J. Milnor, *Local connectivity of Julia sets: expository lectures*. In: "The Mandelbrot Set, Theme and Variations", ed. Tan Lei, 67–116. Cambridge University Press, 2000.
- [Re1] L. Rempe, *A landing theorem for periodic rays of exponential maps*. Proc. Amer. Math. Soc. 134 (2006), no. 9, 2639–2648 (electronic).
- [Re2] L. Rempe, *On nonlanding dynamic rays of exponential maps*. Ann. Acad. Sci. Fenn. Math. 32 (2007), 353–369.
- [RS1] L. Rempe, D. Schleicher, *Combinatorics of bifurcations in exponential parameter space*. In: *Transcendental Dynamics and Complex Analysis*, ed. P. Rippon and G. Stallard, London Mathematical Society Lecture Note Series 348 Cambridge University Press (2008), 317–370.
- [RS2] L. Rempe, D. Schleicher, *Bifurcations in the space of exponential maps*. Invent. Math. 175 (2009), 103–135.
- [RS3] L. Rempe, D. Schleicher, *Bifurcation loci of exponential maps and quadratic polynomials: local connectivity, triviality of fibers, and density of hyperbolicity*. In: *Holomorphic dynamics and renormalization*, 177–196, Fields Inst. Commun., 53, Amer. Math. Soc., Providence, RI, (2008)
- [RvS] L. Rempe, S. Van Strien, *Absence of line fields and Mañé’s theorem for non-recurrent transcendental functions*. Trans. Amer. Math. Soc. 363 (2011), no. 1, 203–228.
- [Ro] P. Roesch, *On local connectivity for the Julia set of rational maps: Newton’s famous example*. Ann. of Math. (2) 168 (2008), no. 1, 127–174.
- [Sc0] D. Schleicher, *On the dynamics of iterated exponential maps*. Habilitation Thesis, TU München (1999).

- [Sc1] D. Schleicher, *On Fibers and Renormalization of Julia Sets and Multibrot Sets*. arXiv:math/9902156.
- [Sc] D. Schleicher, *Attracting dynamics of exponential maps*. Ann. Acad. Sci. Fenn. Math. 28 (2003), 3–34.
- [SZ1] D. Schleicher, J. Zimmer, *Escaping points of exponential maps*. J. London Math. Soc. (2) 67 (2003), no. 2, 380–400.
- [ST] T. Lei, M. Shishikura, *An alternative proof of Mane’s theorem on non-expanding Julia sets*. In: Tan Lei et al., *The Mandelbrot Set, Theme and Variations*, Cambridge Univ. Press (2000).
- [Sl] Z. Slodkowski, *Holomorphic motions and polynomial hulls*. Proc. Amer. Math. Soc. 111 (1991), 347–355.